

Diversity and abundance of nematodes in the sewage of Jodhpur, Rajasthan, India

Razia Sultana¹ & Padma Bohra²

^{1,2}Desert Regional Centre, Zoological Survey of India, Jhalamand, Pali Road, Jodhpur, Rajasthan 342005, India
Email: ¹razia.sultana25@gmail.com (corresponding author),
²bohrapadma@gmail.com

Nematodes are aquatic, semi-aquatic and terricolous organisms in nature and among the most important organisms in decomposer communities. Nematode community structure in polluted water as studied by Beier & Traunspurger (2001) revealed that nematodes can be used as bioindicators of soil health because they are ubiquitous and have diverse feeding behaviours and life strategies (Bongers & Bongers 1998; Neher 2001). Bacterial feeding nematodes have the greatest contribution to the decomposer food web. These decomposer nematode species do not feed directly on organic matter but they graze on microbes and excrete ammonia. Thus, nematodes contribute to nitrogen mineralization. Since nematodes respond rapidly to new resources and the nematode fauna can be analyzed, the structure of the nematode community offers an instrument to assess the condition of the ecosystem.

Date of publication (online): 26 May 2012
Date of publication (print): 26 May 2012
ISSN 0974-7907 (online) | 0974-7893 (print)

Editor: Anonymity requested

Manuscript details:

Ms # o3044
Received 23 December 2011
Final received 06 March 2012
Finally accepted 06 April 2012

Citation: Sultana, R. & P. Bohra (2012). Diversity and abundance of nematodes in the sewage of Jodhpur, Rajasthan, India. *Journal of Threatened Taxa* 4(5): 2614–2616.

Copyright: © Razia Sultana & Padma Bohra 2012. Creative Commons Attribution 3.0 Unported License. JoTT allows unrestricted use of this article in any medium for non-profit purposes, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Acknowledgement: Authors are grateful to Dr.K.Venkataraman for providing research facilities in preparation of this paper.

OPEN ACCESS | FREE DOWNLOAD



Material and methods:

Sediment samples were processed by sieving and decantation and a slightly modified Baermann's funnel technique. Nematodes were fixed in hot formalin. For permanent mounts, specimens were dehydrated by the slow method and mounted in anhydrous glycerin. The individual species population and the total nematode counts were taken. Measurements were made using a drawing tube mounted on a Nikon Eclipse E600 microscope and photographs were taken using a Nikon digital camera DS-Fi1.

Results: A total of 39 species of nematodes were identified from sewage water of various localities of Jodhpur and adjoining areas (Table 1). Five functional nematode trophic groups were identified based on morphological structures and modes of feeding such as herbivores, bacterivores, fungivores, omnivores and predators. In terms of individual abundance, bacterivores (rhabditids) were the most dominant group represented followed by predators, herbivores, omnivores and fungivores. In terms of taxonomic diversity cephalobids were the most abundant group followed by rhabditis/diplogastrids among the bacteriovores. *Mononchoides*, *Mesorhabditis* and *Diplogasteritus* were found the most dominant genera in the sewage water.

Discussion: Nematode frequency, density and diversity vary depending upon ecological and edaphic factors (Sohlenius 1979; Khatoon et al. 2001). The present study revealed a great deal of generic diversity within the nematode community. A total of 39 nematode species were identified, representing bacterivores (14 genera), predators (six genera), omnivores (three genera), herbivores (two genera) and fungivores (one genus). Fifty species of nematodes representing five trophic groups were reported in the sewage waters (Tahseen 2006). Most of the nematode species were microbe grazing which have been reported to regulate the rates of decomposition (Yeates & Coleman 1982). Increased bacterivores diversity shows the increased diversity of microbes thus reflecting the nature and quality of the environment. This present study shows a high nematode density and low species diversity in the sewage water.

Table 1. Diversity of various trophic groups in sewage system of Jodhpur.

List of species	cp value	Trophic group
Tylenchida Thorne, 1949		
Hoplolaimidae Filipjev, 1934	(3)	Herbivore
<i>Helicotylenchus dihystra</i> (Cobb, 1893) Sher, 1961	(3)	Herbivore
<i>Helicotylenchus erythrinae</i> (Zimmerman, 1904) Golden, 1956	(3)	Herbivore
Criconematidae Taylor, 1936		
<i>Hemicriconemoides brachyurus</i> (Loos, 1949) Chitwood & Birchfield, 1957	(3)	Herbivore
Aphelenchida Siddiqi, 1980		
Aphelenchidae Fuchs, 1937		
<i>Aphelenchus avenae</i> Bastian, 1865	(2)	Fungivore
Dorylaimida Pearse, 1942		
Dorylaimidae de Man, 1876		
<i>Mesodorylaimus subtiloides</i> (Paetzold, 1958) Andrassy, 1959	(4)	Omnivore
<i>Mesodorylaimus kauli</i> Baqri & Bohra, 2001	(4)	Omnivore
Aporcelaimidae Heyns, 1965		
<i>Aporcelaimellus heynsi</i> Baqri & Jairajpuri, 1968	(5)	Predator
Qudsianematidae Jairajpuri, 1965		
<i>Ecumenicus monhystera</i> (De Man, 1880) Thorne, 1974	(4)	Omnivore
Nordiidae Jairajpuri & Siddiqi, 1964		
<i>Kochinema farodai</i> Baqri & Bohra, 2001	(4)	Omnivore
Actinolaimidae Thorne, 1939		
<i>Neoactinolaimus rajasthanensis</i> Bohra & Sultana, 2008	(4)	Predator
Mononchida Jairajpuri, 1969		
Mylonchulidae Jairajpuri, 1969		
<i>Mylonchulus brachyurus</i> (Bütschli, 1873)	(4)	Predator
Rhabditida Chitwood, 1933		
Cephalobidae Filipjev, 1934		
<i>Cephalobus cubaensis</i> Steiner, 1935	(2)	Bacterivore
<i>Cephalobus quadrilineatus</i> Eroshenko, 1968	(2)	Bacterivore
<i>Chiloplacus sclerovaginitus</i> Sumenkova & Razzhivin, 1968	(2)	Bacterivore
<i>Heterocephalobus pulcher</i> (Loof, 1964) Andrassy, 1967	(2)	Bacterivore
<i>Acrobeles chelatus</i> Thomas & Allen, 1965	(2)	Bacterivore
<i>Acrobeles geraerti</i> (Rashid et al, 1990) Shahina & De Ley, 1997	(2)	Bacterivore
<i>Acrobeles mariannae</i> Andrassy, 1968	(2)	Bacterivore
<i>Zeldia neoacuta</i> Allen & Noffsinger, 1972	(2)	Bacterivore
<i>Zeldia feria</i> Allen & Noffsinger, 1972	(2)	Bacterivore
Teratocephalidae		
<i>Teratocephalus bisexualis</i> Meyer & Coomans, 1977	(1)	Bacterivore
Panagrolaimidae Thorne, 1937		
<i>Panagrolaimus peruensis</i> (Steiner, 1939) Goodey, 1963	(1)	Bacterivore
<i>Panagrolaimus dendroctoni</i> (Fuchs, 1932) Rühm, 1956	(1)	Bacterivore
Rhabditidae Örley, 1880		
<i>Mesorhabditis anisomorpha</i> (Sudhaus, 1978) Andrassy, 1983	(1)	Bacterivore
<i>Mesorhabditis miotki</i> (Sudhaus, 1978) Andrassy, 1983	(1)	Bacterivore
<i>Teratorhabditis synpapillata</i> Sudhaus, 1985	(1)	Bacterivore
Diploscapteridae Micoletzky, 1922		
<i>Diploscapter coronatus</i> (Cobb, 1893) Cobb, 1913	(1)	Bacterivore
Diplogastridae Micoletzky, 1922		
<i>Butlerius butleri</i> Goodey, 1929	(4)	Predator
<i>Diplogasteritis nudicapitatus</i> (Steiner, 1914) Paramonov, 1952	(4)	Predator
Neodiplogastridae Paramonov, 1952		
<i>Mononchoides longicaudatus</i> (Khera, 1965) Andrassy, 1984	(4)	Predator
Araeolaimida De Coninck & Schuurmans Stekhoven, 1933		
Leptolaimidae Örley, 1880		
<i>Chronogaster daoi</i> Loof, 1964	(2)	Bacterivore
<i>Chronogaster typica</i> (De Man, 1921) De Coninck, 1935	(2)	Bacterivore
Chromadorida Chitwood, 1933		
Cyatholaimidae Filipjev, 1918		
<i>Achromadora micoletzkyi</i> (Stefanski, 1915) Van Der Linde, 1938	(2)	Bacterivore
<i>Achromadora ruricola</i> (de Man, 1880) Micoletzky, 1925	(2)	Bacterivore
Monhysterida Schuurmans Stekhoven & De Coninck, 1933		
Monhysteridae de Man, 1876		
<i>Monhystera africana</i> Andrassy, 1964	(2)	Bacterivore
<i>Monhystera paludicola</i> de Man, 1881	(2)	Bacterivore
Enoplida Baird, 1853		
Prismatolaimidae Micoletzky, 1922		
<i>Prismatolaimus andrassyi</i> Khera & Chaturvedi, 1967	(3)	Bacterivore
<i>Prismatolaimus intermedius</i> (Bütschli, 1873) De Man, 1880	(3)	Bacterivore
Tripylidae De Man, 1846		
<i>Tobrilus longus</i> (Leidy, 1852) Andrassy, 1959	(4)	Predator

REFERENCES

- Beier, S. & W. Traunspurger (2001).** The meiofauna community of two German streams as indicator of pollution. *Journal of Aquatic Ecosystem Stress and Recovery* 8: 387–405.
- Bongers, T. & M. Bongers (1998).** Functional diversity of nematodes. *Applied Soil Ecology* 10: 239–251.
- Khatoon, M., S. Sharma & S.K. Saxena (2001).** Community analysis of predatory nematodes of Rohilkhand division. U.P. *Current Nematology* 12: 11–14
- Neher, D.A. (2001).** Role of nematodes in soil health and their use as indicators. *Journal of Nematology* 33: 161–168.
- Sohlenius, B. (1979).** A carbon budget for nematodes, rotifers and tardigrades in a Swedish coniferous forest sil. *Holarctic Ecology* 2: 30–40.
- Tahseen, Q. (2006).** Taxonomic diversity in sewage nematodes of Aligarh, North India. *International Journal of Nematology* 16: 1–6.
- Yeates, G.W. & D.C. Coleman (1982).** Nematodes in decomposition, pp. 55–80. In: Freckman, D.W. (ed.). *Nematodes in Soil Ecosystems*. University of Texas, Austin TX, USA.

