

## OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.



## Journal of Threatened Taxa

Building evidence for conservation globally

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

### NOTE

**NEW LOCALITY RECORDS OF THE STOUT SAND SNAKE  
*PSAMMOPHIS LONGIFRONS* BOULENGER, 1890 (REPTILIA:  
SQUAMATA: LAMPROPHIIDAE) IN TELANGANA, INDIA**

Avinash C. Visvanathan, Sandeep Anne & Aditya Kesav Kolli

26 November 2017 | Vol. 9 | No. 11 | Pp. 10968–10970

10.11609/jott.3449.9.11.10968–10970



For Focus, Scope, Aims, Policies and Guidelines visit [http://threatenedtaxa.org/About\\_JoTT](http://threatenedtaxa.org/About_JoTT)

For Article Submission Guidelines visit [http://threatenedtaxa.org/Submission\\_Guidelines](http://threatenedtaxa.org/Submission_Guidelines)

For Policies against Scientific Misconduct visit [http://threatenedtaxa.org/JoTT\\_Policy\\_against\\_Scientific\\_Misconduct](http://threatenedtaxa.org/JoTT_Policy_against_Scientific_Misconduct)

For reprints contact <[info@threatenedtaxa.org](mailto:info@threatenedtaxa.org)>

Partner



Publisher/Host





ISSN 0974-7907 (Online)  
ISSN 0974-7893 (Print)

#### OPEN ACCESS



The Stout Sand Snake *Psammophis longifrons* Boulenger, 1890 is a rare, endemic Indian snake recorded from Thane, Maharashtra (Dereckman 1892), Thane, Damanganga, Bulsar, Panch Mahals and Nagpur, Maharashtra and Gujarat (Smith 1943), Surat, Gujarat (Vyas 1987), Amravati, Melghat and Buldana, Maharashtra (Nande & Deshmukh 2007; Joshi 2011), Ujjain, Madhya Pradesh (Ingle 2009), Hoshangabad, Madhya Pradesh (Kumbhar et al. 2012), Navsari and Valsad, Gujarat (Vyas & Patel 2013), Mokhada, Maharashtra (Bansode et al. 2016), Shimoga, Karnataka (Premkumar & Sharma 2017) (Image 1). Here we record the presence of *P. longifrons* in Telangana.

On 26 November 2015, at 11:00hr, J. Prabhakar, a member of the “Friends of Snakes Society”, rescued a *Psammophis longifrons* (Image 2A) that had entered a suburban residence in Boduppal, Hyderabad (Buddha Nagar, Boduppal 17°24'51.6636”N & 78°35'44.4516”E, elevation 498m). This is the first record from the state of Telangana. Another live specimen (Image 2B) was rescued by Aparna Ravi, another member of the same organization, on 22 January 2016, at 11:44hr, while responding to a distress call from Hayathnagar (G.V.R. Colony, Tattiannaram, Hayathnagar 17.36361111 N & 78.60166667 E, 488m).

The specimens were collected, studied and photographed, before their subsequent relocation, close to their home range. Morphometric measurements and pholidosis were measured using standard methods as defined by Whitaker & Captain (2004).

**Specimen 1 (S1):** female, measuring 700mm, snout

## NEW LOCALITY RECORDS OF THE STOUT SAND SNAKE *PSAMMOPHIS LONGIFRONS* BOULENGER, 1890 (REPTILIA: SQUAMATA: LAMPROPHIIDAE) IN TELANGANA, INDIA

Avinash C. Visvanathan<sup>1</sup>, Sandeep Anne<sup>2</sup> & Aditya Kesav Kolli<sup>3</sup>

<sup>1,2,3</sup> Friends of Snakes Society, 34-114/1, Vivekanandapuram, Sainikpuri, Secunderabad, Telangana 500094, India  
<sup>1</sup>avinash@friendsofsnakes.org (corresponding author),  
<sup>2</sup>annesandeep969@gmail.com, <sup>3</sup>adityakesav@gmail.com

to vent length 510mm, tail length 190mm. Dorsal scales (counted one head-length behind the head, at mid-body and at one head length anterior to the vent) were in rows of 17:17:13; ventrals 154; subcaudals 88 (paired); supralabials 8, with 4<sup>th</sup> and 5<sup>th</sup> in contact with the eye; postoculars 2; preocular 1 (grooved); loreal 1 (elongated, horizontally grooved); nasal scales divided into one anterior and two posterior nasals; temporals 2+3; anal scale divided.

**Specimen 2 (S2):** female, measuring 1031mm, snout to vent length 920mm, tail length 111mm (a portion of the tail was missing). Dorsal scales (counted one head-length behind the head, at mid-body and at one head length anterior to the vent) were in rows of 17:17:13; ventrals 165; subcaudals 20 paired (a portion of the tail was missing); supralabials 8, with 4<sup>th</sup> and 5<sup>th</sup> in contact with the eye; postoculars 2; preocular 1 (grooved); loreal 1 (elongated, horizontally grooved); nasal scales divided into one anterior and two posterior nasals; temporals 2+3; anal scale divided.

Scales were smooth and lustrous; dorsum was pale olive and the ventral part was pale yellow. Paraventral

DOI: <http://doi.org/10.11609/jott.3449.9.11.10968-10970>

Editor: Gernot Vogel, Heidelberg, Germany.

Date of publication: 26 November 2017 (online & print)

Manuscript details: Ms # 3449 | Received 18 April 2017 | Final received 02 October 2017 | Finally accepted 01 November 2017

Citation: Visvanathan, A.C., S. Anne & A.K. Kolli (2017). New locality records of the Stout Sand Snake *Psammophis longifrons* Boulenger, 1890 (Reptilia: Squamata: Lamprophiidae) in Telangana, India. *Journal of Threatened Taxa* 9(11): 10968–10970; <http://doi.org/10.11609/jott.3449.9.11.10968-10970>


Copyright: © Visvanathan et al. 2017. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: None.

Competing interests: The authors declare no competing interests.

Acknowledgements: We thank the Telangana Forest Department for permitting us to study the specimens in captivity. We also wish to thank Aparna Ravi for helping us with the morphometric assessment.





**Image 1.** Documented distribution records of *P. longifrons* - 1. Cuddapah, Andhra Pradesh (Type locality); 2. Thane, Maharashtra (Dreckmann 1892; Smith 1943); 3. Damanganga, Maharashtra (Smith 1943); 4. Valsad, Gujarat (Smith 1943; Vyas & Patel 2013); 5. Panch Mahals, Gujarat (Smith 1943); 6. Nagpur, Maharashtra (Smith 1943); 7. Surat, Gujarat (Vyas 1987); 8. Amravati, Maharashtra (Nande & Deshmukh 2007); 9. Melghat, Maharashtra (Nande & Deshmukh 2007); 10. Buldana, Maharashtra (Joshi 2011); 11. Ujjain, Madhya Pradesh (Ingle 2009); 12. Navsari, Gujarat (Vyas & Patel 2013); 13. Shimoga, Karnataka (Premkumar & Sharma 2017); 14. Boduppal, Telangana (present study); 15. Hayathnagar, Telangana (present study)

scales or the outermost dorsal scale rows were bi-coloured—top half resembled the dorsum colouration, while the bottom half was pale yellow, like the ventral scales. Two longitudinal pale yellow stripes, running parallel, were present on the dorsum, from cervical region to the tail tip. Each dorsal scale, between these two stripes, was enveloped by a black border. Broad black margins surrounded the vertebral scales, forming a continuous chain along the back. Black symmetrical markings were present on the parietal, frontal and supraocular scales.

While briefly being kept in captivity before its relocation, it was observed that *P. longifrons* was strictly diurnal, feeding on geckoes, *Hemidactylus* sp. (Image 2C). Prey was actively pursued and subdued before being ingested head first. Both specimens were terrestrial; however, they climbed vegetation with ease. Rescue sites of both individuals were surrounded by dry deciduous thorny vegetation.

Current reports from the state of Telangana, India are mere accidental encounters, due to rapid urbanization and habitat fragmentation. There is an urgency to carry out thorough scientific assessments to identify

distribution ranges and support concrete efforts to conserve this rare species.

## References

- Boulenger, G.A. (1890).** *The Fauna of British India, Including Ceylon and Burma. Reptilia and Batrachia.* Taylor & Francis, London, xviii, 541pp.
- Dreckmann (1892).** Miscellaneous Notes. No V. A Rare Snake. The Journal of the Bombay Natural History Society vii, 406pp.
- Ingle, M. (2009).** First locality record of Stout Sand Snake, *Psammophis longifrons* Boulenger, 1896 from Ujjain District, Madhya Pradesh. *Cobra* 3(2): 12–13.
- Joshi, P.S. (2011).** A preliminary survey on the snakes of Buldhana district, Maharashtra. *Golden Research Thoughts* 1(2): 1–4.
- Nande, R. & S. Deshmukh (2007).** Snakes of Amravati district including Melghat, Maharashtra, with important records of the Indian Egg-eater, Montane Trinket Snake and Indian Smooth Snake. *Zoos' Print Journal* 22(12): 2920–2924; <http://doi.org/10.11609/JoTT.ZP.1653.2920-4>
- Premkumar, S. & V. Sharma (2017).** Geographic distribution: *Psammophis longifrons* (Stocky Sand Racer). *Herpetological Review* 48(1): 131.
- Smith, M.A. (1943).** *The Fauna of British India, Ceylon and Burma, Including the Whole of the Indochinese Subregion* Vol. III. Serpentes, Taylor and Francis, London.
- Vyas, R. (1987).** Collection of Stout Sand Snake *Psammophis longifrons*. *Hamadryad* 12(2): 7–8.
- Vyas, R. & H. Patel (2013).** Notes on distribution and natural



Image 2. *Psammophis longifrons*: A - Full body (S1); B - Lateral close-up of the head (S1); C - S2 feeding on *Hemidactylus* sp.

history of *Psammophis Longifrons* Boulenger, 1896 (Serpentes: Psammophiidae: Psammophiinae) in Gujarat, India. *Russian Journal of Herpetology* 20(3): 217–222.

Whitaker, R. & A. Captain (2004). *Snakes of India - The Field Guide*. Draco Books, Chennai, 480pp.







OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

November 2017 | Vol. 9 | No. 11 | Pages: 10865–10984

Date of Publication: 26 November 2017 (Online & Print)

DOI: 10.11609/jott.2017.9.11.10865-10984

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

#### Communication

**Flies matter: a study of the diversity of Diptera families (Insecta: Diptera) of Mumbai Metropolitan Region, Maharashtra, India, and notes on their ecological roles**

-- Aniruddha H. Dhamorikar, Pp. 10865–10879

#### Short Communications

**Small carnivores of the montane forests of Eravikulam National Park in the Western Ghats, India**

-- S. Nikhil & P.O. Nameer, Pp. 10880–10885

**Distribution and population of Himalayan Marmot *Marmota himalayana* (Hodgson, 1841) (Mammalia: Rodentia: Sciuridae) in Leh-Ladakh, Jammu & Kashmir, India**

-- Vipin Chaudhary, R.S. Tripathi, Surjeet Singh & M.S. Raghuvanshi, Pp. 10886–10891

**First record of Bourret's Horseshoe Bat *Rhinolophus paradoxolophus* (Mammalia: Chiroptera: Rhinolophidae) from Myanmar with a review of the taxonomy, distribution and ecology of the species**

-- Sai Sein Lin Oo, Du Sar No, Lucia Nang Seng, Ngwe Lwin, Malcolm Pearch & Paul J.J. Bates, Pp. 10892–10898

**A first record of the Smallfin Gulper Shark *Centrophorus moluccensis* Bleeker, 1860 (Chondrichthyes: Squaliformes: Centrophoridae) from the Andaman & Nicobar waters, Indian EEZ**

-- H.D. Pradeep, Swapnil S. Shirke, M. Nashad & Monalisha Devi Sukham, Pp. 10899–10903

**Taxonomic revision of the genus *Atmetonychus* (Coleoptera: Curculionidae: Entiminae) from the Indian subcontinent**

-- G. Mahendiran & V.V. Ramamurthy, Pp. 10904–10908

**A new species of dewflower *Murdannia sanjappae* (Commelinaceae) from Andaman Islands, India**

-- Mudavath Chennakesavulu Naik & Boyina Ravi Prasad Rao, Pp. 10909–10913

**First records of two Ginger Lily *Hedychium* (Zingiberaceae) species from the Western Ghats, India**

-- Sinjumol Thomas, Susai John Britto & Bince Mani, Pp. 10914–10919

**An annotated checklist of microbes associated with bamboo in the Indian subcontinent**

-- O.K. Remadevi, P. Sharada & H.C. Nagaveni, Pp. 10920–10947

#### Notes

**Roadkill records of Lowland Tapir *Tapirus terrestris* (Mammalia: Perissodactyla: Tapiridae) between kilometers 06 and 76 of highway BR-163, state of Pará, Brazil**

-- Marco A. de Freitas, Rodrigo C. Printes, Eric K. Motoyama, Assor E. Fucks & Diogo Veríssimo, Pp. 10948–10952

**Population size, herd structure and sex ratio of the Blackbuck *Antelope Cervicapra* (Mammalia: Cetartiodactyla: Bovidae) in a human dominated area in Odisha, India**

-- Subrat Debata, Pp. 10953–10955

**Recovery of Musk Deer *Moschus chrysogaster* Hodgson, 1839 (Artiodactyla: Moschidae) in Sakteng Wildlife Sanctuary, Bhutan**  
-- Sonam Tobgay, Thinley Wangdi & Kumbu Dorji, Pp. 10956–10958

**First record of the Asiatic Brush-tailed Porcupine *Atherurus macrourus* Linnaeus, 1758 (Mammalia: Rodentia: Hystricidae) from western Bhutan**

-- Tashi Dhendup & Rinzin Dorji, Pp. 10959–10960

**The Vulnerable Indian Skimmer *Rynchops albicollis* Swainson, 1838 (Aves: Charadriiformes: Laridae) breeding in Odisha, eastern India**

-- Subrat Debata, Tuhinansu Kar, Kedar Kumar Swain & Himanshu Shekhar Palei, Pp. 10961–10963

**On the occurrence of Black Baza *Aviceda leuphotes* Dumont, 1820 (Aves: Falconiformes: Accipitridae) in the Guptaeswar forests of the Eastern Ghats, Odisha, India**

-- Swetashree Purohit, Manoj V. Nair & Sharat Kumar Palita, Pp. 10964–10967

**New locality records of the Stout Sand Snake *Psammodon longifrons* Boulenger, 1890 (Reptilia: Squamata: Lamprophiidae) in Telangana, India**

-- Avinash C. Visvanathan, Sandeep Anne & Aditya Kesav Kolli, Pp. 10968–10970

**A note on the distribution of two highly threatened butterflies in Sri Lanka (Lepidoptera: Lycaenidae: *Spindasis greeni* and *Rapala lankana*), with a report on the range extension of *S. greeni***  
-- Tharaka Sudesh Priyadarshana, Ishara Harshajith Wijewardhane & Mithila Karunaratna, Pp. 10971–10973

**A new record of grass *Ottolochloa* (Poaceae) to the Eastern Ghats, India**

-- Midigesi Anil Kumar, P. Anjaneyulu & Boyina Ravi Prasad Rao, Pp. 10974–10976

**An extended distribution of Natesh's Cape-pondweed *Aponogeton nateshii* (Aponogetonaceae), a new record to the state of Goa**

-- Rutuja Rajendra Kolte, Anup Satish Deshpande, Prabha Muraleedharan Pillai & Shrirang Ramchandra Yadav, Pp. 10977–10979

**Detection of *Artyfechinostomum sufrartyfex* - a zoonotic parasite from the Small Indian Mongoose *Herpestes auropunctatus* (Mammalia: Carnivora: Herpestidae) in Jammu & Kashmir, India**

-- Sanku Borkataki, Pankaj Goswami, Rajesh Katoch, Sahil Kumar & Pratiksha Raghuvanshi, Pp. 10980–10982

#### Book Review

**Requisite for long term studies in ecology**

-- S. Suresh Ramanan, Pp. 10983–10984

**zoo!**  
Z Ü R I C H

**WILD  
ZOO**  
Threatened Taxa