

Total number of printed pages-7

3 (Sem-6/CBCS) MAT HC 2

2022

**MATHEMATICS**

(Honours)

Paper : MAT-HC-6026

**(Partial Differential Equations)**

Full Marks : 60

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer **any seven** :  $1 \times 7 = 7$

(i) The equation of the form

$P_p + Q_q = \mathbb{R}$  is known as

(a) Charpit's equation

(b) Lagrange's equation

(c) Bernoulli's equation

(d) Clairaut's equation

(Choose the correct answer)

Contd.

(ii) How many minimum no. of independent variables does a partial differential equation require?

(iii) Find the degree and order of the equation

$$\frac{\partial^3 z}{\partial x^3} + \left( \frac{\partial^3 z}{\partial x \partial y^2} \right)^2 + \frac{\partial z}{\partial y} = \sin(x + 2y)$$

(iv) Which method can be used for finding the complete solution of a non-linear partial differential equation of first order

- (a) Jacobi method
- (b) Charpit's method
- (c) Both (a) and (b)
- (d) None of the above

(Choose the correct answer)

(v) State True Or False :

The equation

$$u_{xx} + u_{yy} + u_{zz} = 0$$

is an Hyperbolic equation.

(vi) Fill in the blanks :

$$\left( \frac{\partial z}{\partial x} \right)^2 + 2 \frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial y^2} + z = 0$$

is a \_\_\_\_\_ order partial differential equation.

(vii) The characteristic equation of  $yu_x + xu_y = u$  is

(a)  $\frac{dx}{x} = \frac{dy}{y} = \frac{du}{u}$

(b)  $\frac{dx}{y} = \frac{dy}{x} = \frac{du}{u}$

(c)  $\frac{dx}{u} = \frac{dy}{x} = \frac{du}{y}$

(d) None of the above

(Choose the correct answer)

(viii) State True Or False

$xu_x + yu_y = u^2 + x^2$  is a semi-linear partial differential equation.

(ix) Fill in the blanks :

A solution  $z = z(x, y)$  when interpreted as a surface in 3-dimensional space is called \_\_\_\_\_.

(x) The partial differential equation is elliptical if

(a)  $B^2 - 4AC > 0$

(b)  $B^2 - 4AC \geq 0$

(c)  $B^2 - 4AC \leq 0$

(d)  $B^2 - 4AC < 0$

(Choose the correct answer)



2. Answer **any four** :  $2 \times 4 = 8$

(i) Define quasi-linear partial differential equation and give *one* example.

(ii) Show that a family of spheres

$(x-a)^2 + (y-b)^2 = r^2$  satisfies the partial differential equation

$$z^2(p^2 + q^2 + 1) = r^2$$

(iii) Eliminate the constants  $a$  and  $b$  from  $z = (x+a)(y+b)$ .

(iv) Determine whether the given equation is hyperbolic, parabolic or elliptic

$$u_{xx} - 2u_{yy} = 0.$$

(v) Solve the differential equation  $p + q = 1$ .

(vi) Explain the essential features of the "Method of separation of variables".

(vii) Mention when Charpit's method is used. Name a disadvantage of Charpit's method.

(viii) What is the classification of the equation

$$u_{xx} - 4u_{xy} + 4u_{yy} = e^y$$

3. Solve **any three** :  $5 \times 3 = 15$

(i) Form a partial differential equation by eliminating arbitrary functions  $f$  and  $F$  from  $y = f(x-at) + F(x+at)$ .

(ii) Solve

$$y^2 p - xyq = x(z - 2y)$$

(iii) Find the integral surface of the linear partial differential equation

$$x(y^2 + z)p - y(x^2 + z)q = (x^2 - y^2)z$$

which contains the straight line

$$x + y = 0, \quad z = 1.$$

(iv) Find the solution of the equation  $z = pq$  which passes through the parabola

$$x = 0, \quad y^2 = z.$$

(v) Find a complete integral of the equation

$$x^2 p^2 + y^2 q^2 = 1.$$

(vi) Reduce the equation  $yu_x + u_y = x$  to canonical form and obtain the general solution.



(vii) Apply the method of separation of variables  $u(x, y) = f(x)g(y)$  to solve the equation  $u_x + u = u_y$ ,

$$u(x, 0) = 4e^{-3x}.$$

(viii) Determine the general solution of

$$4u_{xx} + 5u_{xy} + u_{yy} + u_x + u_y = 2.$$

4. Answer **any three** :  $10 \times 3 = 30$

(i) Solve  $(p^2 + q^2)y - qz = 0$  by Jacobi method.

(ii) Solve  $z^2 = pqxy$  by Charpit's method.

(iii) Find the general solution of the differential equation

$$x^2 \frac{\partial z}{\partial x} + y^2 \frac{\partial z}{\partial y} = (x + y)z$$

(iv) Solve

$$(mz - ny)p + (nx - lz)q = ly - mx$$

(v) Use  $v = \ln u$  and  $v = f(x) + g(y)$  to solve the equation

$$x^2 u_x^2 + y^2 u_y^2 = u^2.$$

(vi) Find the solution of the equation

$$z = \frac{1}{2}(p^2 + q^2) + (p - x)(q - y)$$

which passes through the  $x$  axis.

(vii) Find the canonical form of the equation

$$y^2 u_{xx} - x^2 u_{yy} = 0.$$

(viii) Classify the second order linear partial differential equation with example.